

AD\_\_\_\_\_

Award Number: W81XWH-05-1-0117

TITLE: Prospective study of ALS mortality among World War II, Korea, and Vietnam Veterans

PRINCIPAL INVESTIGATOR: Alberto Ascherio, M.D., Dr.Ph.

CONTRACTING ORGANIZATION: Harvard University  
Cambridge, MA 02138-3846

REPORT DATE: January 2006

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. <b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b>					
1. REPORT DATE (DD-MM-YYYY) 01-01-2006		2. REPORT TYPE Annual		3. DATES COVERED (From - To) 31 DEC 2004 - 30 DEC 2005	
4. TITLE AND SUBTITLE Prospective study of ALS mortality among World War II, Korea, and Vietnam				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-05-1-0117	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Alberto Ascherio, M.D., Dr.Ph.  E-mail: aascheri@hsph.harvard.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Harvard University Cambridge, MA 02138-3846				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT We prospectively assessed the association between ALS mortality and self-reported military service in the American Cancer Society's Cancer Prevention Study II (CPS II) cohort, a cohort that includes over 500,000 men surveyed by questionnaire in 1982. ALS mortality was assessed via linkage with the National Death Index. Our original analyses(3) included ALS deaths (n=280) up to 1998. We have now extended the follow-up to 2002 (ALS deaths =513). Mantel-Haenszel relative risks (RR) adjusted for age and smoking were calculated and Cox proportional hazards models stratified on single year of age were used when adjusting for additional variables. Men who served in the military (69% of our population) had a significantly increased ALS mortality (RR=1.5) compared to those who did not serve. The increase in ALS mortality was similar among men who served in the Army or National Guard (RR=1.6), Navy (RR=1.7), or Air Force (RR=1.5). This increased risk of ALS appeared largely independent of the branch of service, years of service, and war periods during service. The project will continue as planned. We will extend the follow-up and examine the possible role of vitamin E supplements in prevention.					
15. SUBJECT TERMS Amyotrophic lateral sclerosis, military service, vitamin E, epidemiology					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT  UU	18. NUMBER OF PAGES  7	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

## Table of Contents

Cover.....	1
SF 298.....	2
Introduction.....	4
Body.....	4
Key Research Accomplishments.....	4
Reportable Outcomes.....	4
Conclusions.....	4
References.....	4
Appendices.....	5

**Introduction:** Two recent studies suggested that the risk of Amyotrophic Lateral Sclerosis (ALS) is increased among Gulf War veterans.(1, 2) Initial work of ours suggested that military service prior to the Gulf War was also associated with an increase in risk for ALS.(3) Our purpose here was to extend our previous investigation by extending follow-up an extra 4 years to see whether the previous associations persisted.

**Body:** We prospectively assessed the association between ALS mortality and self-reported military service in the American Cancer Society's Cancer Prevention Study II (CPS II) cohort, a cohort that includes over 500,000 men surveyed by questionnaire in 1982. ALS mortality was assessed via linkage with the National Death Index. Our original analyses(3) included ALS deaths (n=280) up to 1998. We have now extended the follow-up to 2002 (ALS deaths =513). Mantel-Haenszel relative risks (RR) adjusted for age and smoking were calculated and Cox proportional hazards models stratified on single year of age were used when adjusting for additional variables. Men who served in the military (69% of our population) had a significantly increased ALS mortality (RR=1.5; 95% CI: 1.2-2.0; p = 0.0009) compared to those who did not serve. The increase in ALS mortality was similar among men who served in the Army or National Guard (RR=1.6), Navy (RR=1.7), or Air Force (RR=1.5). The increased risk of ALS did not vary much by years of service (figure 1), the number of major conflicts during service (figure 2), nor service specifically in World War 2, the Korean war, or the Vietnam war (figure3).

**Key Research Accomplishments:**

- We have completed the analyses of the extended follow-up of military service and risk of ALS death in the ACS CPS-II cohort through 2002.

**Reportable Outcomes:** Presentation of this work at the 22<sup>nd</sup> Neurotoxicology Conference in Raleigh Durham, September, 2005.

**Conclusions:** In extended analyses with almost twice as many ALS cases as our previous work, we confirmed an increased risk of ALS mortality among US military veterans who completed their period of service before the first Gulf War. This increased risk of ALS appeared largely independent of the branch of service, years of service, and war periods during service. The project will continue as planned. We will extend the follow-up and examine the possible role of vitamin E supplements in prevention.


**References:**

1. R. W. Haley, *Neurology* **61**, 750 (2003).
2. R. D. Horner *et al.*, *Neurology* **61**, 742 (2003).
3. M. G. Weisskopf *et al.*, *Neurology* **64**, 32 (Jan 11, 2005).

**Figure 1.** Relative risk of ALS by years of service, adjusted for smoking, education, alcohol intake, and vitamin E use.



**Figure 2.** Relative risk of ALS by number of major war periods during service, adjusted for smoking, education, alcohol intake, and vitamin E use.



**Figure 3.** Relative risk of ALS by service during major war period, adjusted for smoking, education, alcohol intake, and vitamin E use.

